arXiv:1201.1989v2 [astro-ph.CO] 16 Jan 2012

Volume Title
*ASP Conference Series, Vol. **Volume Number***
Author
© ***Copyright Year** Astronomical Society of the Pacific*

Andromeda and its Satellites - a Kinematic Perspective

Michelle L. M. Collins,[1] R. Mike Rich,[2] and Scott C. Chapman[3]

[1]*MPIA, Königstuhl 17, Heidelberg, 69117, Germany*

[2]*UCLA, Physics and Astronomy Building, Los Angeles, CA 90095-1547, USA*

[3]*Institute of Astronomy, Madingley Rise, Cambridge, CB2 1ST, UK*

Abstract. Using spectroscopic data taken with Keck II DEIMOS by the Z-PAndAS team in M31, I present a comparison of the disc and satellite systems of M31 with those of the Galaxy. I discuss observed discrepancies between the masses and scale radii of M31 dwarf spheroidal galaxies of a given luminosity with those of the Milky Way. I also also present an analysis of the newly discovered M31 thick disc, which is measured to be hotter, more extended and thicker than that seen in the Milky Way.

1. The Thick Disc in Andromeda

Thick stellar discs are believed to represent an earlier epoch of galactic evolution than thin stellar discs. Understanding the properties of these structures is fundamental to our understanding of the evolution of large spiral galaxies. Until recently, no such component was detected in Andromeda, despite the belief that such structures are ubiquitous amongst spiral galaxies (Dalcanton & Bernstein 2002). We present a detection of a thick disc in the Andromeda galaxy (Collins et al. 2011a), isolating the component by identifying a significant population of stars that have velocities that lag behind the thin stellar disc, and are inconsistent with the Andromedean halo.

We have been able to measure the scale lengths of the discs. We measure $h_{r,thin}$= 7.3 kpc and $h_{r,thick}$= 8.0 kpc. Comparing these values with thick disc systems in external galaxies (Yoachim & Dalcanton 2006, YD06) we estimate the scale height of both components. We measure $z_{0,thin}$= 1.1 kpc and $z_{0,thick}$= 2.8 kpc. We co-add their spectra to ascertain the average metallicity of each component via the measurement of the equivalent widths of the Ca II triplet (Fig. 1). The thick disc has an intermediate metallicity of [Fe/H]= -1.0, more metal poor than the thin disc ([Fe/H]= -0.7) and more metal rich than the halo ([Fe/H]= -1.3).

2. The Dwarf Spheroidals of Andromeda

Dwarf spheroidal galaxies (dSphs) represent the smallest scales on which we are able to detect dark matter. 27 dSphs have been identified in the M31 halo, and while the relationship between the size and luminosity of these objects is consistent with the MW dSphs (Brasseur et al. 2011), there are a number that are more extended than their MW counterparts (McConnachie & Irwin 2006; Richardson et al. 2011, Fig. 1).



Figure 1. **Left:** Composite spectra for the thin (top), thick disc (middle) and halo (bottom) populations in M31. These are chemically distinct, with the thick disc having a metallicity of [Fe/H]= −1.0. **Centre:** Velocity dispersion vs. half light radius for MW (red) M31 (blue) and isolated (light blue) dSphs. The best fit mass profiles of Walker et al. (2009) are overlaid. A number of M31 dSphs have lower velocity dispersions than expected. **Right:** Ratio of the measured : expected velocity dispersion from W09 NFW profile for Local Group dSphs vs. surface brightness. The M31 outliers are seen in the lower left corner. A tidal track from Penarrubia et al. (2008) representing a dSph that has been stripped of 90 % of its mass is overplotted, and matches the observed trend.

Work by Walker et al. (2009) demonstrated that the MW dSphs were consistent with having formed in halos with identical dark matter profiles, suggesting that the evolution of the dark matter within these objects is the same, regardless of the size of the luminous component. Our group has been measuring the kinematics of the M31 dSphs (Chapman et al. 2005, 2007; Collins et al. 2010, 2011b). We have discovered that these objects are not consistent with the MW population, as a number of them have lower velocity dispersions for a given half light radius when compared to the MW. These outliers are also typically the more extended M31 dSphs (Fig. 1), consistent with the effects of increased tidal stress from a more massive M31 disc (Peñarrubia et al. 2008, 2010). Recent work by the SPLASH collaboration (Tollerud et al. 2011) also note some outliers in their survey, but conclude that overall, the difference between the masses of MW and M31 dSphs are not statistically significant.

References

Brasseur, C. M., et al. 2011, ApJ, 743, 179
Chapman, S. C., et al. 2005, ApJ, 632, L87
— 2007, ApJ, 662, L79
Collins, M. L. M., et al. 2010, MNRAS, 407, 2411
— 2011a, MNRAS, 413, 1548
— 2011b, MNRAS, 417, 1170
Dalcanton, J. J., & Bernstein, R. A. 2002, AJ, 124, 1328
McConnachie, A. W., & Irwin, M. J. 2006, MNRAS, 365, 1263
Peñarrubia, J., Navarro, J. F., & McConnachie, A. W. 2008, ApJ, 673, 226
Peñarrubia, J., et al. 2010, MNRAS, 406, 1290
Richardson, J. C., et al. 2011, ApJ, 732, 76
Tollerud, E. J., et al. 2011, ArXiv e-prints. `1112.1067`
Yoachim, P., & Dalcanton, J. J. 2006, AJ, 131, 226